

Register @ Times Times Of Oman >> International News >> Where fear resides

Letter to Editor Tuesday, January 25, 2005



[Start a discussion on this topic](#)

Reader's Forum

PDF Archives

Healthy Times By Roxanne Khamsi

Edit your Profile The New York Times

NEWS

NEW YORK — A woman who cannot recognise fear in people's faces is causing neuroscientists to rethink theories of how our brains read emotions.

Scientists have been testing the 38-year-old woman for more than a decade. She has a rare disease that has damaged both sides of her amygdala, the almond-shaped part of the brain that is known from imaging studies to be involved in recognising facial expressions.

The woman, known as SM, finds it impossible to tell from facial expressions when another person is afraid, although she has no problem recognising other emotions, such as happiness, sadness and anger.

Originally, researchers thought this meant that different emotions are processed by distinct neural circuits in the brain. But new studies with SM, published in Nature recently, suggest quite a different explanation.

Neuroscientist Ralph Adolphs of the California Institute of Technology in Pasadena and his colleagues showed SM photographs of various different facial expressions and asked her what emotion she thought the people were feeling.

Researchers have done this many times before. But this time, to find the part of the faces on which SM was focusing, the team used a "bubble test" in which only part of the face is revealed at a time.

The researchers were intrigued to find that SM totally avoided looking at people's eyes. She discerned her information simply from looking around the nose and mouth. This was generally enough for her to identify emotions such as happiness or anger,

News in this Section

- ▶ [Where fear resides](#)
- ['The Aviator' takes flight](#)
- [US hostage begs for life](#)
- [Over 300 killed in Indian stampede](#)
- [Freed Chinese hostages arrive home](#)
- [India celebrates Republic Day amidst blasts](#)
- [Palestine, Israel resume political contact](#)

Today's Poll

Tourism will boost tsunami-hit nations. Do you agree?

Yes

No

Not Sure

HomePage
 Local
 ▶ International
 Business
 Sports
 ICE Times
 Education

ePaper New

OPINIONS

Editorial
 Viewer Reviews
 Articles
 Columns

BUSINESS

Oman Economy
 Tenders
 Oil Prices
 Stock Prices
 Currency Rates

INFORMATION

Prayer Timings
 Weather
 Horoscope
 Events
 Movies in Town
 Reviews
 Audio/Video
 TV/Radio Schedule
 Flight Schedule
 Shipping Schedule
 Duty Pharmacies

Click to Read

Healthy Times

Hospitals

Clinics

Police

E-GREETINGS

where features such as a smile, or bared teeth, are important.

But wide eyes are a particularly important component of a fearful expression. Because SM was only looking at the nose and mouth, she did not notice the eyes and concluded that the person was feeling neutral.

SPECIAL

Royal Decrees

Law

Classifieds

"First you have to look at the eyes, and then the brain has to make use of that information to figure out it's fear," explains Adolphs. When the researchers simply told SM to look at people's eyes, she did, and her ability to distinguish fear dramatically improved.

Advertise @ Times

She needed constant reminding, as otherwise she stopped looking at the eyes. But this ability to regain lost function simply through receiving the correct instructions is extremely unusual in patients with brain damage. It suggests that SM doesn't have a problem processing the visual cues at all.

Instead, says Adolphs, the area she has damaged may tell us where and how to look. "It's a radical rethink of amygdala function," he says. If our brains simply took in all the information around us, we would be completely overwhelmed, he points out. "What's needed is selectivity of some kind, to decide what's important," he says. Adolphs believes that the amygdala is a critical component of such a mechanism.

The team suggests that this new understanding could be used to help develop treatments for people with disorders such as autism, who find it difficult to read emotions.

Autism sufferers are known to focus abnormally on certain facial features. Giving them direct reminders of how to look at other people's faces could help them to improve their perception of social cues.