

[Register for Updates](#)[About This Site](#)[Editorial Board](#)[Help/FAQ](#)[Comments](#)

BioEd Online

BIOLOGY TEACHER RESOURCES
FROM BAYLOR COLLEGE OF MEDICINE[Home](#) [Presentations](#) [Slides](#) [Sets](#) [Teacher Resources](#) [Biology News](#) [Prof Dev](#) [Discussions](#) [Hot Topics](#)

BIOLOGY NEWS

[Disaster conference](#)

The World Conference on Disaster Reduction, hosted by the United Nations in Kobe, Japan, from 18-22 January, comes at a good time: in the wake of December's tsunami, researchers and policy makers are both keen to do as much as possible to reduce the death

[Titan reveals methane rain and rocks of water](#)

A week's analysis confirms Earth-like features on saturnian moon.

[Booze boosts brainpower](#)

Moderate alcohol consumption protects women from cognitive decline.

[Prions discovered in unexpected organs](#)

Immune system helps BSE proteins spread through the body.

[Y chromosome reveals hidden sequence](#)

Extra genes may determine men's height and health.

[Cuttlefish win mates with transvestite antics](#)

Sneaky male mimics don a female disguise in an instant.

[More Biology News](#)

RECOMMENDED DISCUSSIONS

[tell me one beneficial mutation example - if there is](#)

in [Topica Miscellaneous](#)

[Teachable moment](#)

in [In the Classroom](#)

[MAT Program at Bard College needs Biology Students](#)

January 5, 2005

Terror shows only in the eyes

Knowing where to look is key to recognizing others' emotions.

by Roxanne Khamssi
news@nature.com

A woman who cannot recognize fear in people's faces is causing neuroscientists to rethink theories of how our brains read emotions.

Scientists have been testing the 38-year-old woman for more than a decade. She has a rare disease that has damaged both sides of her amygdala, the almond-shaped part of the brain that is known from imaging studies to be involved in recognizing facial expressions.

The woman, known as SM, finds it very difficult to tell from facial expressions when another person is afraid, although she has no problem recognizing other emotions, such as happiness, sadness and anger.

Originally, researchers thought this meant that different emotions are processed by distinct neural circuits in the brain. But new studies with SM, published in this week's Nature, suggest quite a different explanation.

A closer look

“This is a radical rethink of amygdala function.”

Ralph Adolphs

Neuroscientist at the California Institute of Technology, Pasadena

Neuroscientist Ralph Adolphs of the California Institute of Technology in Pasadena and his colleagues showed SM photographs of various different facial expressions and asked her what emotion she thought the people were feeling.

Researchers have done this many times before. But this time, to find the part of the faces on which SM was focusing, the team used a 'bubble test' in which only part of the face is revealed at a time.

The researchers were intrigued to find that SM totally avoided looking at people's eyes. She discerned her information simply from looking around the nose and mouth.

This was generally enough for her to identify emotions such as happiness or anger, where features such as a smile, or bared teeth, are important.

But wide eyes are a particularly important component of a fearful expression. Because

RELATED SITE

- [Fear and the amygdala](#)

[Register for updates when new articles are posted](#)

[Discuss this article](#)

in **The Teaching Profession**

Joint Project

in **Science Education Partnerships**

Neurology and Sports

in **Topica Miscellaneous**

Uterine Transplant

in **The BioEd Online Community**

Discussions Home

SM was only looking at the nose and mouth, she did not notice the eyes and concluded that the person was feeling neutral.

"First you have to look at the eyes, and then the brain has to make use of that information to figure out it's fear," explains Adolphs.

Rescue effort

When the researchers simply told SM to look at people's eyes, she did, and her ability to distinguish fear dramatically improved.

She needed constant reminding, as otherwise she stopped looking at the eyes. But this ability to regain lost function simply through receiving the correct instructions is extremely unusual in patients with brain damage. It suggests that SM doesn't have a problem processing the visual cues at all.

Instead, says Adolphs, the area she has damaged may tell us where and how to look. "It's a radical rethink of amygdala function," he says.

If our brains simply took in all the information around us, we would be completely overwhelmed, he points out. "What's needed is selectivity of some kind, to decide what's important," he says.

Adolphs believes that the amygdala is a critical component of such a mechanism.


Potential treatments

The team suggests that this new understanding could be used to help develop treatments for people with disorders such as autism, who find it difficult to read emotions.

Autism sufferers are known to focus abnormally on certain facial features. Giving them direct reminders of how to look at other people's faces could help them to improve their perception of social cues.

1 Adolphs R., et al. Nature, 433. 68 - 72 (2005).

Article Copyright © 2005 [MacMillan Publishers Ltd.](#) All rights reserved. This material may not be published, broadcast, rewritten, or redistributed.





BioEd Online is a partnership between Baylor College of Medicine and Texas A&M University, and is funded by grants from the Houston Endowment, Inc.; Howard Hughes Medical Institute; the US Department of Education; and the Science Education Partnership Award program of the National Center for Research Resources, National Institutes of Health.

© 2004–2005 Baylor College of Medicine. All Rights Reserved.

Terms of Use and Privacy Policy

fruit fly image © 2001 Dennis Kunkel Microscopy, Inc.